


Panel discussion
Posterior malleolus

Haraguchi
I, II, III

Bartonicek/Rammelt
1, 2, 3, 4

Mac

Conclusions None of the reviewed PMF classifications has been able to establish itself decisively in the literature. Most of the classifications are insufficient in terms of a derivable treatment algorithm or a prognosis with regard to outcome. However, as the Bartonicek/Rammelt classification has the greatest potential due to its treatment algorithm, its reliability in combination with consistent predictive values, its usage in clinical practice and research appears advisable.



Day 0

60 y male
Slipped on a ramp
No co-morbidities
(however, weight 105 kg, BMI 38)

Reduced in ER




Day 2



1500. Ib1 3D animation




1500. Ib1 3D animation



1500. Ib1 3D animation

Day 2




Day 7



Rounding up

Scott's parabola: the rise and fall of a surgical technique



J W Scott consultant gynaecologist, Poole Hospital NHS Trust, Poole, Dorset BH15 2JB

Article



Open Reduction and Internal Fixation of the Posterior Malleolus Fragment in Ankle Fractures **Improves the Patient-Rated Outcome**: A Systematic Review

Rainer Christoph Miksch, MD ¹, Viktoria Herterich, MD ¹, Alexej Barg, MD, PhD², Wolfgang Böcker, MD¹, Hans Polzer, MD ¹, and Sebastian Felix Baumbach, MD ¹

European Journal of Orthopaedic Surgery & Traumatology (2023) 33:2525–2532
<https://doi.org/10.1007/s00590-022-03455-0>

ORIGINAL ARTICLE

Open fixation of the posterior malleolus **increases the morbidity** of trimalleolar ankle fracture fixation

Shreyaas Aravindan¹ · Nicholas J. Tucker² · Parker J. Prusick¹ · Cyril Mauffrey^{1,2} · Joshua A. Parry^{1,2} 

Received: 29 August 2022 / Accepted: 1 December 2022 / Published online: 29 December 2022
© The Author(s), under exclusive licence to Springer-Verlag France SAS, part of Springer Nature 2022

BE AWARE
Confirmation bias


and SPIN



Tim White, Edinburgh

Guildal lecture 2023

Bartonicek/Rammelt
1, 2, 3, 4



**Assess not only the posterior malleolus
but also, the entire fracture/ligament injury
and the WHOLE PATIENT**

RESEARCH

Open Access



Trans-Achilles percutaneous fixation technique for posterior malleolus fracture: a cadaveric study

Hilal Yağar^{1*}, Fatih Çiçek², Faruk Gazi Ceranoğlu², Hüseyin Karadağ³ and Selim Çınaroğlu²

